

DATE:	03/06/08
NAME:	Mold Solutions
Address:	4458 Augusta Rd Ste 4C
City/ST/Zip:	Lexington, SC 29073
Telephone:	803-996-0116

Mailing Address for samples:

The EnviroScreening Lab
 3665 E. Bay Drive #204
 PMB 428
 Largo, Florida 33771

MOLD SCREENING REPORT

Client NAME:	Dr Bledsoe
Address:	41 Boardwalk Lane
City/ST/ZIP:	Lexington, SC 29072
Telephone:	

DATE Sample #1:	02/28/08
DATE Sample #2:	02/28/08
DATE Sample #3:	
DATE Sample #4:	

	LOCATION/SOURCE	Identification	COUNT
Sample #1	#5 Over door Mstr to LR:	Penicillium sp.	8 cfu
swab		Curvularia sp.	1 cfu
		Total colonies:	9 cfu
Sample #2	#6 Supply front BR:	Cladosporium sp.	9 cfu
swab		Penicillium sp.	11 cfu
		Rhizopus sp.	2 cfu
		Total colonies:	22 cfu

* **cfu**: colony forming units per square inch.
 * **TNTC**: too numerous to count per square inch (>200cfu)
 * for **Hyphal Fragments**, see following Page

Please see RANGE BROCHURE for General Organism Information and Disclaimers.

YOUR REPORT SHOULD INCLUDE THREE PAGES as follows:

1. Screening Report
2. Acceptable Ranges Graph
3. Reference/Disclaimer Sheet

ACCEPTABLE RANGES GRAPH



EnviroScreening, LLC
State of Louisiana LLC
1.877.251.2296

This Reference Sheet has been produced by and is the exclusive property of www.EnviroScreening.com designed for the express purpose of providing analytical information regarding the interpretation of your mold screening kit results.

Specific questions should be directed to
TechSupport@EnviroScreening.com

Should I be concerned about the level of contamination I've found in my report? Maybe. This screening kit provides an inexpensive alternative to other types of air sampling. It is, by all standards, extremely accurate in determining the viability of mold spores in our indoor environments. By testing settled-out spores using this methodology, you will have an accurate representation of what is going on throughout the environment. However, your level of concern should be based on your sensitivity to the types of mold(s) found in your specific environment rather than the amount of contamination. Some people are extremely sensitive to mold and even the lowest level found (Expected/Normal) might prove to be an allergen for that person. At the other extreme, some people have virtually no sensitivity to mold(s) and, for them, mold at higher levels is not a serious concern. In my book, "Mold-Free Construction", I discuss the importance of screening for all types of mold and I talk about methods by which you can have a mold-free, allergy-free environment. When you work with a NORMI Certified IAQ/Mold Professional, you can be sure that the suggestions made for your specific environment will provide for you and your family a mold-free environment for years to come. When you have adopted those recommendations, you need not be concerned!

D. Douglas Hoffman
 National Marketing Director
 Executive Director for www.NORMI.org
 Credentials Available at www.EnviroScreening.com

Contamination LEVELS CHART/GRAPH

<u>SWAB/PETRI DISH ANALYSIS</u>		
GENERAL Interpretation	Single ¹	Mixed ²
Probable Colonization (high contamination)	>100	>300
	100cfu	300cfu
Potential Colonization (moderate contamination)	50cfu	150cfu
	Potential Growth (limited contamination)	15cfu
Expected/Normal (minor contamination)		

<u>TAPE LIFT/BULK SAMPLING ANALYSIS</u>	
Relative Abundance of Mold Spores or Items Identified	
<10 spores/item, total lift	Expected / Normal
10-25: spores/item, total lift	Potential Growth
10-25: spores/item, per field	Potential Colonization
>25: spores/item, per field	Probable Colonization
NOTE: Air-O-Cell samples, unless otherwise noted, all counts obtained are raw counts of total air trace. No calculations performed.	

¹ Single is a single identified organism
² Mixed is a combined of identified organisms

FOR MORE INFORMATION
www.EnviroScreening.com
www.MoldFreeConstruction.com

RANGES GUIDELINE DISCLAIMERS

- 1) It should be noted that mold and bacteria is found throughout our environment and identifying mold in your environment does not necessarily mean that you have a mold problem.
- 2) Currently there are no Federal or State recognized standards, critical or threshold exposure limits to aeroallergens.
- 3) Because everyone's immune system is unique, individual exposure response to aeroallergens will vary. The purpose of these guidelines is to help the user determine the need and value of advise received from a Certified IAQ Specialist.
- 4) These guidelines are unique only to these test methodologies and have been determined to be reasonable by the manufacturer of the test kits (EnviroScreening.com) in conjunction with NORMI (National Organization of Remediators & Mold Inspectors) Counts on the report are estimates based on the density of growth.
- 5) Test results only apply to the area from which the sample was taken, and the description of the ranges only signify the increasing amount of germs present.
- 6) This methodology should only be considered a screening, and more detailed analysis may be necessary. For more information, discuss additional testing options with your Certified IAQ Specialist.
- 7) **These screening methodologies are mold screenings only and therefore NOT intended for legal / lawsuit purposes. Any attempt to use them for legal purposes will be considered hostile and a distortion of their intended use which is to provide a reasonably priced screening alternative to more expensive testing procedures. Should legal documentation be required we recommend that the client or customer consider utilizing the services of a Licensed Industrial Hygenist who can control testing procedures.**

REFERENCE/DISCLAIMER SHEET

Many pages have been written regarding mold, mildew and the deleterious effects these have or may have on construction materials and our health. For more information, please review our website at www.EnviroScreening.com and read the book entitled **Mold-Free Construction™** which is available at www.MoldFreeConstruction.com.

In an effort to help you sort through the information, we are providing this reference sheet along with information regarding ranges and the potential contamination that certain ranges may represent. Please read all the footnotes so you will have a clear understanding of what this screening test does and does not represent. Drawing conclusions from information that a screening test provides is best done by a certified IAQ Specialist.

In most cases, and certainly where extreme remediation measures are considered, this screening test should be understood as simply being a “snap-shot” of what MAY be going on in other areas of your indoor environment. Settled-out spores can often be representative of sampling a larger area but the correlation is still unclear between what you find in a square inch sampling and what you might find in other areas of the environment.

The material found in this brochure has been reviewed by the NORMI (National Organization of Remediators and Mold Inspectors) and is provided in agreement with their standards. Any specific questions may be directed to:
Info@EnviroScreening.com

www.EnviroScreening.com

www.MoldFreeConstruction.com

REFERENCE COMMENT SHEET—*There are many different types of bacteria that can be found throughout our living environment. These may inhabit dust particulate, soil and non-treated water. Examples: species of Bacillus, actinomycete and nonfermentative gram-negative rods. Infections from most of these isolates are not common, however, occurrence of infection will vary based on species isolated.*

Acromonium sp. is another common fungus that is found in our environment, which has been reported to cause such ailments as mycetomas (swelling of tissues), keratitis (cornea), and other mycoses.

Alternaria sp. is commonly found in our environment to include such places as soil, plants and decaying vegetation. Has been reported to cause keratitis (cornea), skin and nasal infections.

Aspergillus is found throughout our environment and disease caused by this fungus is relatively uncommon. However, there are species of this genus that can cause both invasive and non-invasive diseases, most notably with immunosuppressed hosts. Allergic response to inhalation of the aspergilli has also been known to occur.

Bipolaris sp. is commonly isolated in the laboratory from environmental samples, it has been reported to cause sinusitis and allergies in some cases.

Chaetomium sp. is a common laboratory “contaminant” naturally found in the soil, plants, or decaying vegetation. Infection is not common.

Chrysosporium is a fungus that rarely causes disease. However, cases of endocarditis and osteomyelitis has been reported. Several species are keratinophilic and can cause nail infections.

Cladosporium is another common mold isolated in the laboratory. It is naturally found in soil and in association with plant material. Infection is also uncommon; however, infections of the sinuses, endocarditis, keratomycosis (cornea) and subcutaneous lesions have been noted.

Curvularia is a commonly isolated environmental mold and some species has been noted to cause disease in man.

Epicoccum is commonly isolated in the laboratory based on geographic location and has been associated with allergies.

Fusarium sp. is a common laboratory saprophyte commonly found in soil and decaying vegetation. Fusarium has been reported to cause keratitis (cornea).

Geotrichum sp. is another filamentous fungi that is considered a common laboratory contaminant, infections due to this organism is rare in man.

Hyphal Fragments: are fragments/broken pieces of molds that are not complete structures and cannot, therefore, be fully identified.

Monilia sp. is a common laboratory filamentous fungi that can be found in samples containing soil. This organism is a common laboratory contaminant and infection is not common.

Paecilomyces sp. is a fast growing, common environmental fungus. Some species have been noted in cases of endocarditis, sinusitis, nephritis and infections in dogs.

Penicillium is a fungus that is abundant in man’s environment and is a common laboratory contaminant. Penicillium has also been known to cause a number human infections such as keratitis (eye / cornea), penicilliosis, otomycosis (ear), and rarely, deep infections.

Rhizopus sp. is a common laboratory contaminant. Infection is not common, however, cases of mucormycosis (various types of fungal infections) has been noted.

Trichoderma sp. is a common filamentous fungi. Infections from this organism is not common but various ailments has been reported.

Stachybotrys chartarum has been well publicized for being a potential agent in causing severe respiratory problems in young children and persons with immunosuppressed conditions. It is commonly referred to as “toxic mold” & “black mold”.

Sterile Hyphae: are molds/fungi grown in culture that has not produced spores and cannot, therefore, be accurately identified.

Syncephalastrum sp. is a common environmental mold which is fast spreading in culture, nonseptate and could be mistaken for *Aspergillus niger*. A case of cutaneous infection has been reported.

Yeast are seen in nature from sources like soil, fruits, milk, plants, and feces of normal humans. Infections vary based on the species and sources identified.

NOTE: *There is scientific evidence showing that persons with immunosuppressed conditions run the highest risk of infection due to environmental factors such as bacteria and fungal elements. Please see more at www.EnviroScreening.com*

REFERENCES/DISCLAIMERS

†This information is contained in medical mycology and microbiology textbooks such as: *Medical Mycology* by K.J. Kwon-Chung & John E. Bennett, *Diagnostic Microbiology* by Koneman, Bailey & Scott’s *Diagnostic Microbiology* by Ellen Jo Baron, Lancer R. Peterson & Sydney M. Finegold and *Medical Mycology* by Martha E. Kern & Kathleen S. Blevins. **More information:** www.cdc.gov/nceh/airpollution/mold/stachy.htm

* **This information should only be used as general information for identification purposes only and not for diagnostic purposes.**

* **It should be noted that mold & bacteria are found throughout our environment and identifying mold in your environment does not necessarily mean that you have a mold problem.**

* Cultures submitted may have colony counts estimated based on the density of growth in the culture.

* Because acceptable counts may vary from laboratory to laboratory and are not well established, it is important to take cultures before treatment of facility and again after treatment of facility in order to ensure adequate reduction of colony forming units. Additionally, collection techniques should be performed the same way each time a sample is taken

* **This methodology should be considered a screening technique only. A more extensive analysis may be necessary. Please ask you IAQ Specialist for interpretation of these results.**

Recommendation: Air purification and mold remediation companies can help reduce the amount of particulate matter in the air, including the mold spore counts. (See your local Certified IAQ Specialist for more information and recommendations regarding air purification equipment and remediation options)

Glenn Moore Jr. BS. MT. Supv.
Florida license # SU 30654
Glen@EnviroScreening.com